

A note on the span of Hadamard products of vectors

Hajime Tanaka

*Division of Mathematics, Graduate School of Information Sciences, Tohoku University
6-3-09 Aramaki-Aza-Aoba, Aoba-ku, Sendai 980-8579, Japan
E-mail: htanaka@math.is.tohoku.ac.jp*

Abstract

We give a new proof of Theorem 6 in [L. Qiu and X. Zhan, On the span of Hadamard products of vectors, Linear Algebra Appl. 422 (2007) 304–307].

AMS classification: 15A03; 15A04; 15A57

Keywords: Hadamard product; Range

The *Hadamard product* $A \circ B$ of two $m \times n$ complex matrices $A = (A_{ij}), B = (B_{ij})$ is defined by their entry-wise multiplication, i.e., $(A \circ B)_{ij} = A_{ij}B_{ij}$. Qiu and Zhan [3] studied the range of the Hadamard product of $n \times n$ complex matrices. In this note, we give a new proof of the following theorem:

Theorem 1 ([3, Theorem 6]). *Let $B_1, B_2, \dots, B_k$ be $n \times n$ complex matrices. Then*

$$\text{span}\{(B_1x_1) \circ \dots \circ (B_kx_k) | x_1, \dots, x_k \in \mathbb{C}^n\} = \text{range}((B_1B_1^*) \circ \dots \circ (B_kB_k^*)),$$

where $*$ means conjugate transpose.

Proof. Let e_i be the vector in $\mathbb{C}^n$ with a 1 in the i th coordinate and 0 in all other coordinates. Then for $1 \leq i \leq n$ the i th column of $(B_1B_1^*) \circ \dots \circ (B_kB_k^*)$ is

$$((B_1B_1^*) \circ \dots \circ (B_kB_k^*))e_i = (B_1B_1^*e_i) \circ \dots \circ (B_kB_k^*e_i).$$

So

$$\text{range}((B_1B_1^*) \circ \dots \circ (B_kB_k^*)) \subseteq \text{span}\{(B_1x_1) \circ \dots \circ (B_kx_k) | x_1, \dots, x_k \in \mathbb{C}^n\}.$$

We show

$$\text{span}\{(B_1x_1) \circ \dots \circ (B_kx_k) | x_1, \dots, x_k \in \mathbb{C}^n\} \subseteq \text{range}((B_1B_1^*) \circ \dots \circ (B_kB_k^*)). \quad (1)$$

Pick any $x_1, x_2, \dots, x_k \in \mathbb{C}^n$. Let E be the orthogonal projection onto $\text{range}((B_1B_1^*) \circ \dots \circ (B_kB_k^*))^\perp$. Then for any $y \in \mathbb{C}^n$ we have

$$\begin{aligned} \langle (B_1x_1) \circ \dots \circ (B_kx_k), Ey \rangle &= \sum_{i=1}^n \langle B_1x_1, e_i \rangle \dots \langle B_kx_k, e_i \rangle \overline{\langle Ey, e_i \rangle} \\ &= \sum_{i=1}^n \langle x_1 \otimes \dots \otimes x_k \otimes \bar{y}, (B_1^*e_i) \otimes \dots \otimes (B_k^*e_i) \otimes \bar{E}e_i \rangle \\ &= \langle x_1 \otimes \dots \otimes x_k \otimes \bar{y}, \sum_{i=1}^n (B_1^*e_i) \otimes \dots \otimes (B_k^*e_i) \otimes \bar{E}e_i \rangle, \end{aligned}$$

where $\bar{}$ means complex conjugate. On the other hand, using $E^* = E^2 = E$ we have

$$\| \sum_{i=1}^n (B_1^*e_i) \otimes \dots \otimes (B_k^*e_i) \otimes \bar{E}e_i \|^2 = \sum_{i,j=1}^n \langle B_1^*e_i, B_1^*e_j \rangle \dots \langle B_k^*e_i, B_k^*e_j \rangle \langle \bar{E}e_i, \bar{E}e_j \rangle$$

$$\begin{aligned}
&= \sum_{i,j=1}^n (B_1 B_1^*)_{ji} \cdots (B_k B_k^*)_{ji} E_{ij} \\
&= \sum_{i,j=1}^n E_{ij} ((B_1 B_1^*) \circ \cdots \circ (B_k B_k^*))_{ji} \\
&= \text{trace}(E((B_1 B_1^*) \circ \cdots \circ (B_k B_k^*))) \\
&= 0.
\end{aligned}$$

So

$$\sum_{i=1}^n (B_1^* e_i) \otimes \cdots \otimes (B_k^* e_i) \otimes \overline{E} e_i = 0,$$

and we find

$$\langle (B_1 x_1) \circ \cdots \circ (B_k x_k), E y \rangle = 0.$$

Since y is arbitrary, we conclude

$$(B_1 x_1) \circ \cdots \circ (B_k x_k) \in \text{range}((B_1 B_1^*) \circ \cdots \circ (B_k B_k^*)).$$

We have now shown (1) and the result follows. $\square$

Corollary 2 ([3, Theorem 5]). *Let $A_1, A_2, \dots, A_k$ be $n \times n$ positive semidefinite matrices. Then*

$$\text{span}\{(A_1 x_1) \circ \cdots \circ (A_k x_k) | x_1, \dots, x_k \in \mathbb{C}^n\} = \text{range}(A_1 \circ \cdots \circ A_k).$$

Proof. Set $B_i = A_i^{\frac{1}{2}}$ ($1 \leq i \leq k$) in Theorem 1 and recall $\text{range}(A_i) = \text{range}(B_i)$ ($1 \leq i \leq k$). $\square$

Corollary 3 ([3, Theorem 4]). *Let $A_1, A_2, \dots, A_k$ be $n \times n$ positive semidefinite matrices. Then*

$$\text{span}\{(A_1 x) \circ \cdots \circ (A_k x) | x \in \mathbb{C}^n\} = \text{range}(A_1 \circ \cdots \circ A_k).$$

Proof. For $1 \leq i \leq n$ the i th column of $A_1 \circ \cdots \circ A_k$ is

$$(A_1 \circ \cdots \circ A_k) e_i = (A_1 e_i) \circ \cdots \circ (A_k e_i).$$

So by Corollary 2 we have

$$\begin{aligned}
\text{range}(A_1 \circ \cdots \circ A_k) &\subseteq \text{span}\{(A_1 x) \circ \cdots \circ (A_k x) | x \in \mathbb{C}^n\} \\
&\subseteq \text{span}\{(A_1 x_1) \circ \cdots \circ (A_k x_k) | x_1, \dots, x_k \in \mathbb{C}^n\} \\
&= \text{range}(A_1 \circ \cdots \circ A_k),
\end{aligned}$$

as desired. $\square$

Remark. Obviously Corollary 2 in turn implies Theorem 1 (cf. [3, Theorem 6]). Qiu and Zhan [3] proved Corollary 2 first and got the other theorems. Their proof makes full use of positive semidefiniteness, and the key lemma there [3, Lemma 3] needs a slight analytic argument [4, p. 15]. The idea of taking tensor products in this note is from [2]. See also [1, Section 2.8].

Acknowledgments. The author would like to thank Akihiro Munemasa and Hiroki Tamura for comments and discussions. Thanks are also due to Xingzhi Zhan for carefully reading the manuscript.

References

- [1] E. Bannai and T. Ito, Algebraic combinatorics I: Association schemes, Benjamin/Cummings, Menlo Park, CA, 1984.
- [2] P. J. Cameron, J.-M. Goethals and J. J. Seidel, The Krein condition, spherical designs, Norton algebras and permutation groups, *Nederl. Akad. Wetensch. Indag. Math.* 40 (1978) 196–206.
- [3] L. Qiu and X. Zhan, On the span of Hadamard products of vectors, *Linear Algebra Appl.* 422 (2007) 304–307.
- [4] X. Zhan, Matrix inequalities, *Lecture Notes in Mathematics*, 1790, Springer-Verlag, Berlin, 2002.